

Presentation 7.3b: Multiple linear regression

Murray Logan

09 Aug 2016

Preparations

PACKAGES

```
library(ggplot2)
library(car)
library(GGally)
library(rstan)
library(brms)
library(coda)
library(dplyr)
library(gridExtra)
```

DATA

www.flutterbys.com.au/stats/downloads/data/loyn.csv

www.flutterbys.com.au/stats/downloads/data/paruelo.csv

Section 1

Theory

Multiple Linear Regression

ADDITIVE MODEL

growth = intercept + temperature + nitrogen

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_j x_{ij} + \epsilon_i$$

OR

$$y_i = \beta_0 + \sum_{j=1:n} \beta_j x_{ji} + \epsilon_i$$

Multiple Linear Regression

ADDITIVE MODEL

growth = intercept + temperature + nitrogen

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_j x_{ij} + \epsilon_i$$

- effect of one predictor holding the other(s) constant

Multiple Linear Regression

ADDITIVE MODEL

growth = intercept + temperature + nitrogen

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \dots + \beta_j x_{ij} + \epsilon_i$$

Y	X1	X2
3	22.7	0.9
2.5	23.7	0.5
6	25.7	0.6
5.5	29.1	0.7
9	22	0.8
8.6	29	1.3
12	29.4	1

Multiple Linear Regression


MULTIPLICATIVE MODEL

growth = intercept + temp + nitro + temp × nitro

$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \beta_3 x_{i1} x_{i2} + \dots + \epsilon_i$$

Assumptions

- normality, homogeneity of variance, linearity
- (multi)collinearity



Multiple Linear Regression

VARIANCE INFLATION

$$\text{var.inf} = \frac{1}{1 - R^2}$$

Collinear when $\text{var.inf} \geq 5$

Some prefer > 3

Section 2

Worked Examples

Worked examples

```
loyn <- read.csv('../data/loyn.csv', strip.white=T)
head(loyn)
```

	ABUND	AREA	YR.	ISOL	DIST	LDIST	GRAZE	ALT
1	5.3	0.1	1968	39	39	2	160	
2	2.0	0.5	1920	234	234	5	60	
3	1.5	0.5	1900	104	311	5	140	
4	17.1	1.0	1966	66	66	3	160	
5	13.8	1.0	1918	246	246	5	140	
6	14.1	1.0	1965	234	285	3	130	

Worked Examples

Question: what effects do fragmentation variables have on the abundance of forest birds

Linear model:

$$\text{Abund}_i \sim \mathcal{N}(\mu, \sigma^2)$$

$$\mu = \beta_0 + \sum_{j=1:n}^N \beta_j x_{ji}$$

$$\beta_0, \beta_j \sim \mathcal{N}(0, 1000)$$

$$\sigma \sim \text{Cauchy}(0, 5)$$

Worked Examples

```
paruelo <- read.csv('../data/paruelo.csv', strip.white=T)
head(paruelo)
```

	C3	LAT	LONG	MAP	MAT	JJAMAP	DJFMAP
1	0.65	46.40	119.55	199	12.4	0.12	0.45
2	0.65	47.32	114.27	469	7.5	0.24	0.29
3	0.76	45.78	110.78	536	7.2	0.24	0.20
4	0.75	43.95	101.87	476	8.2	0.35	0.15
5	0.33	46.90	102.82	484	4.8	0.40	0.14
6	0.03	38.87	99.38	623	12.0	0.40	0.11

Worked Examples

Question: what effects do fragmentation geographical variables have on the abundance of C3 grasses

Linear model:

$$\sqrt{C3_i} \sim \mathcal{N}(\mu, \sigma^2)$$

$$\mu = \beta_0 + \sum_{j=1:n}^N \beta_j X_{ji}$$

$$\beta_0, \beta_j \sim \mathcal{N}(0, 1000)$$

$$\sigma \sim \text{Cauchy}(0, 5)$$